JAZZY (0-40 mm)

FRANCELINE x CUPIDO



Building on potential

Added value of the variety

Excellent taste, texture and cooking quality.
Very nice skin quality.
Variety with a very high tuberset.
Variety suitable for baby potato production.
Low susceptibility to common scab.



Characteristics

Shape of tuber	L	
Tubers/plant	20	
Av. Tuber length (Ø 50-60 mm)	9,8 cm	
Maturity (9=early, 4=late)	7,6	
Colour of skin	Y	
Colour of flesh	6,9	
Suitability for cooking	8 A	

A B C D

Soil adaptability

	Sandy	Light	Heavy
Yield (Relative)	67	68	82
DM (%)	17,6	18,9	19,4
UWW	317	343	354
Tuber size (Ø mm)	37	38	39

Ro1	2
Ro2/3	1
Ro5	1
Pa2	2
Pa3	1
TRV	7
Moptop virus	9

1 = very susceptible to 9 = resistant

wz 1 (D1)	10
wz 2 (G1)	4
wz 6 (O1)	5
wz 18 (T1)	6

3 = very susceptible to 10 = immune

Climate adaptability

Drought tolerance	7
Secondary growth	7

Resistance / Tolerance

Foliar late blight	3
Tuber late blight	3
Early blight	5
Y-virus	6
Yntn-virus	7
Leafroll virus	
Black dot	9
Silver scurf	9
Metribuzin pre-emergence	9
Metribuzin post-emergence	3

1 = very susceptible to 9 = resistant

Storability

Dormancy* (2=short; 8=long)	2,2
Pressure spots	Rather susceptible
Fusarium (9=resistant, 1=very susceptible)	4
Frying/crisping colour in October*	4,3
Frying/crisping colour in February*	3,5
Frying/crisping colour in May*	3,3

**Potatoes stored at 8°C without sprout inhibitors*

Frying/crisping scale : 8= very good, 3=bad

Marketable yield

Bruising index (0=good-50=bad)	5
Hollow heart (%)	0
IRS (%)	0
Washability	8
Common scab	7
Powdery scab	8
Mechanical Damage	7

1 = very susceptible to 9 = resistant

ROBUST MARKETABLE YIELD

Meijer Potato will not be responsible for possible negative effects originating from this advice.

02/11/2026

JAZZY (0-40 mm)

Planting

Before planting keep the seed quiet; plant them out of cold storage.
Plant at a normal depth.
Growing in bed system gives a real plus to optimise baby production results.
Ask your sales representative for futher advice
Treat the seeds against rhizoctonia and silver scurf.

Size	75 cm	90 cm	Plant/ha
25/30	15 cm	13 cm	88900
30/40	20 cm	17 cm	66700
40/50	26 cm	22 cm	51300
30/50	23 cm	19 cm	58000
50/55	32 cm	27 cm	41700

Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.
To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, Ca)

(N) Nitrogen:	Average needs for an early variety	150	kg/ha	*Netherlands based advice*
(P2O5) Phosphate:	High needs for an early variety	160	kg/ha	
(K2O) Potassium:	Low needs for an early variety	250	kg/ha	



(N) Nitrogen:	100%
(P2O5) Phosphate:	100%
(K2O) Potassium:	100%

In order to increase the tuber set, Phosphate application in the ridge is advised.

Crop protection

Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.
Variety without blight resistant gene, do not neglect blight control.
Quick tuber growing at the end of the cycle, kill in time.

Harvest and storage

Harvest as soon as skin set is done, approx. 18 days after haulm killing.
Short dormancy; after harvest and wound healing cool down quickly,
keep the temperature stable and start with sprout control.
Start sprouting inhibitor application as soon as potatoes are dry.

Storage:
2 °C
6 months

Key advice to succeed this variety:

Before planting keep the seed quiet; plant them out of cold storage.
Growing in bed system gives a real plus to optimise baby production results.
In order to increase the tuber set, Phosphate application in the ridge is advised.
Harvest as soon as skin set is done, approx. 18 days after haulm killing.

Meijer Potato will not be responsible for possible negative effects originating from this advice.

02/11/2026